

The environmental setting of Epipalaeolithic aggregation site Kharaneh IV

Article

Accepted Version

Jones, M. D., Maher, L. A., Macdonald, D. A., Ryan, C., Rambeau, C., Black, S. and Richter, T. (2016) The environmental setting of Epipalaeolithic aggregation site Kharaneh IV. *Quaternary International*, 396. pp. 95-104. ISSN 1040-6182 doi: <https://doi.org/10.1016/j.quaint.2015.08.092> Available at <https://centaur.reading.ac.uk/44811/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1016/j.quaint.2015.08.092>

Publisher: Elsevier

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

The environmental setting of Epipalaeolithic aggregation site Kharaneh IV

Matthew D. Jones^{1*}, Lisa A. Maher², Danielle A. Macdonald^{3,4}, Conor Ryan⁵, Claire
Rambeau⁶, Stuart Black⁷, and Tobias Richter⁸

1. School of Geography, University of Nottingham, University Park, Nottingham, NG7 2RD.
UK
2. Department of Anthropology, University of California, Berkeley, USA
3. Department of Archaeological Science, University of Bradford, UK
4. Department of Anthropology, University of Tulsa, Harwell Hall, 800 S Tucker Drive, Tulsa
OK, 74104
5. Golder Associates (UK) Ltd, Tadcaster, North Yorkshire, UK
6. Institut für Geo- und Umwelt naturwissenschaften – Geologie, University of Freiburg,
Germany
7. Department of Archaeology, School of Archaeology, Geography and Environmental
Science, University of Reading, UK
8. Department of Cross-Cultural and Regional, University of Copenhagen, Copenhagen,
Denmark

*Corresponding author: matthew.jones@nottingham.ac.uk

Abstract

The archaeological site of Kharaneh IV in Jordan's Azraq Basin, and its relatively near neighbour Jilat 6 show evidence of sustained occupation of substantial size through the Early to Middle Epipalaeolithic (c. 24,000 – 15,000 cal BP). Here we review the geomorphological evidence for the environmental setting in which Kharaneh IV was established. The on-site stratigraphy is clearly differentiated from surrounding sediments, marked visually as well as by higher magnetic susceptibility values. Dating and analysis of off-site sediments show that a significant wetland existed at the site prior to and during early site occupation (~ 23,000 – 19,000 BP). This may explain why such a substantial site existed at this location. This wetland dating to the Last Glacial Maximum also provides important information on the palaeoenvironments and potential palaeoclimatic scenarios for today's eastern Jordanian desert, from where such evidence is scarce.

Keywords: Epipalaeolithic, Jordan, Last Glacial Maximum, Wetland, Azraq

1. Introduction and Background

There is much contemporary interest in people's relationships with their natural environment and how resources can be sustainably maintained given changing climates, population sizes, and per capita demands (e.g. Al-Juaidi et al., 2014; Berndtsson et al., 2014). Today people are increasingly vulnerable to risk associated with a changing climate and a finite resource base (e.g. IPCC, 2014). Arguably these issues were also critical for prehistoric societies, although for hunter-gatherers their ability to move around the landscape represented a highly flexible strategy through which climatic change could be effectively mitigated, as long as population levels remained relatively low. In the wider Levant region people's adaptation and mitigation strategies to a changing climate during the transition from the last glacial period into the Holocene interglacial have been widely discussed in relation to the beginnings of agriculture (e.g. Rosen, 2007; Blockley and Pinhasi, 2011; Maher et al., 2011a; Rosen and Rivera-Collazo, 2012). Yet our understanding of how the Levant experienced this global transition in climate is still somewhat unclear (e.g. Robinson et al., 2006; Enzel et al., 2008) and relies on palaeoclimate datasets mainly from the west of the region. To improve our ability to test hypotheses about people's reactions to climatic and environmental change, or about their influence on climate and local environments (e.g. Ruddiman, 2015; Ramsey et al., in press), improved spatial and temporal resolution of our palaeoenvironmental and archaeological records is required (Maher et al., 2011a).

The Azraq Basin of eastern Jordan has long been the focus of archaeological excavation and associated environmental investigations documenting a long history of human occupation dating back to the Lower Palaeolithic (e.g. Field, 1960; Copeland and Hours 1989; Rollefson et al., 1997; Betts, 1998; Garrard and Byrd, 2013). The latest set of excavations in the basin includes work by the Epipalaeolithic Foragers in Azraq Project (EFAP; e.g. Maher et al., 2011b, Maher et al. 2012; Richter et al., 2013; Maher et al., this volume) and this paper reports the results of geomorphological investigations around the site of Kharaneh IV, placing the site into its wider palaeoenvironmental context.

1.1 Kharaneh IV

The Early to Middle Epipalaeolithic site of Kharaneh IV (KHIV) is an important Late Pleistocene site in the Eastern Levant. Recent excavations at KHIV, building on the initial work of M. Muheisen (e.g. 1988), have shown the site to be of great archaeological interest. The high density of artefacts, given a relatively short occupation history (19,830-18,600 cal. years BP; Richter et al., 2013), as well as the thickness of archaeological deposits, large size of the site (22,000 m²), and the presence of very early hut structures (Maher et al., 2012; this

volume), are all rare for Epipalaeolithic sites and suggest frequent re-use of KHIV by hunter-gatherer groups.

The site is located approximately 40km west of the Azraq Oasis (Fig. 1) at an elevation of ~640masl, lying on a sedimentary terrace of pale, cream-coloured silts, in the Wadi Kharaneh, just south of the Islamic castle of the same name. The local topography (Fig. 2) shows the site is the highpoint on the floor of the greater Wadi Kharaneh (Fig. 3); it sits at the confluence of two minor wadis with a general gradient of about 0.3m per 100m to the east, towards the central oasis.

The sediments around the site have been described very briefly before as part of regional reviews (Garrard et al., 1985; Besancon et al. 1989) but before EFAP were not dated or systematically surveyed to link KHIV into the wider landscape. Here we describe such work, providing a geomorphological background to the establishment of KHIV and adding to the palaeoenvironmental reconstruction of the local environment. In combination with faunal data (Martin et al, 2010, Jones, 2012) and ongoing archaeobotanical analysis, this geomorphological data contributes to our understanding of why this particular locality was selected for settlement and why people returned to the same place on the landscape for c. 1000 years (see also Maher, in press). In addition, this work provides more information for an emerging picture of environmental change within the wider Azraq Basin through the late Quaternary (e.g. Jones and Richter, 2011; Cordova et al., 2013; Ames et al., 2014) that improves our understanding of regional environmental and climatic change throughout the Pleistocene and Holocene.

2. Methodology

2.1 Mapping and sediment logging

The topography of the site and the surrounding area was mapped in high-resolution using a ProMark3 differential GPS system, with survey data fixed to the local site grid. In total, 1076 data points were used to create a local contour map of the site and the immediate surrounding area. Six off-site sections were dug into wadi terraces and were visually described and surveyed into the site grid. In addition, a 9m x 1m 'GeoTrench' was dug into the edge of the site itself. Careful surveying of all sections to the site grid allowed these off-site sections to be directly compared to the excavation areas on-site (see Maher et al., this volume for details of these). Of particular interest to this study are the deep sounding in Areas A (excavation square AS42) and B (R/S2/60) and a deep sounding between the two main excavation areas (AZ51), all of which were excavated into the archaeologically sterile units underlying the site.

2. 2 Age-estimates

A number of dating methods have been used to try and constrain the age of the stratigraphy, both on- and off-site, at KHIV. The methodologies for both Optically Stimulated Luminescence (OSL) and U-series approaches are outlined here.

OSL samples were taken in opaque tubes, sealed at both ends, from both off- and on-site sediments (detailed sampling locations are detailed later in the manuscript). On return from the field age estimates were obtained at the University of Gloucestershire Luminescence Dating Laboratory. All samples were opened and prepared under controlled laboratory illumination and to isolate material potentially exposed to daylight during sampling, sediment located within 20 mm of each tube-end was removed. The remaining sample was dried, and then subjected to acid and alkaline digestion to remove carbonate and organic components respectively. Fine silt sized quartz was extracted by sample sedimentation in acetone and feldspars and amorphous silica were then removed from this fraction through acid digestion (Jackson *et al.*, 1976; Berger *et al.*, 1980). Following addition of 10% HCl to remove acid soluble fluorides, grains degraded to $<5\text{ }\mu\text{m}$ as a result of acid treatment were removed by acetone sedimentation. Up to 12 aliquots (ca. 1.5 mg) were then mounted on aluminium discs for Equivalent Dose (D_e) evaluation. D_e values were quantified using a single-aliquot regenerative-dose (SAR) protocol (Murray and Wintle 2000; 2003) measuring the natural signal of a single aliquot and then regenerating that aliquot's signal by using known laboratory doses to enable calibration. For each aliquot, 5 different regenerative doses were administered so as to image dose response. D_e values for each aliquot were then interpolated, and associated counting and fitting errors calculated, by way of exponential plus linear regression. Weighted (geometric) mean D_e values were calculated, given sufficient mass, from 12 aliquots using the central age model outlined by Galbraith *et al.* (1999) and are quoted at 1σ confidence (Table 1). Lithogenic Dose Rate (D_r) values were defined through measurement of U, Th and K radionuclide concentration and conversion of these quantities into α , β and γ D_r values (Table 1). Cosmogenic D_r values were calculated on the basis of sample depth, geographical position and matrix density (Prescott and Hutton, 1994). Ages reported in Table 1 provide an estimate of sediment burial period based on mean D_e and D_r values and their associated analytical uncertainties.

A U-Series age was also obtained from carbonate nodules (e.g. Rowe & Maher 2000) found near the site (see sedimentary descriptions below for details). U/Th data was produced using a Perkin Elmer ELAN 6000 Inductively Coupled Plasma Mass Spectrometer at the University of Reading (e.g. Black *et al.* 2011; Rambeau *et al.* 2011). U/Th dating is based on the

measurement of ^{230}Th produced by the radioactive decay of ^{234}U , once the latter was preferentially incorporated into the newly precipitated sediment. The U/Th technique provides accurate dating only if 1) the system remains chemically closed after deposition, and 2) either if no initial Th is present within the system at time of precipitation, or the amount of additional Th (e.g., as brought in by detrital contamination) can be calculated and corrected for.

Collected samples were composed of dense, micritic carbonate and showed no sign of weathering internally, minimising the likelihood of dating problems due to open-system behaviour. As an attempt to correct for detrital contamination (that would add detrital thorium to the sediment dated, leading to calculation of ages that are too old) an isochron age was also calculated (e.g. Candy et al. 2004, 2005). Both individual and isochron ages were calculated using the program ISOPLOT© V.2.49 (Ludwig 2001), which also provides a statistical assessment of the validity of the calculated best-fit isochron age, by evaluating its relationship to the dataset (Mean Square of Weighted Deviates [MSWD], probability of fit). These statistics are crucial in estimating the accuracy of the calculated isochron age (Candy et al. 2004, 2005 and references therein); a high MSWD (value >1) indicates analytical or geological problems and ages that are potentially more complex..

2.3 Sedimentology

A series of standard sedimentary analyses were undertaken on a set of 52 samples to further quantify the visual sedimentary descriptions; 11 samples from Area A, 24 from Area B and 17 off-site samples (including those from the GeoTrench).

Loss on Ignition analysis was undertaken using standard procedures (e.g. Heiri et al., 2001). Volume-specific Magnetic Susceptibility analysis was undertaken using a Bartington MS2B Dual Frequency Magnetic Susceptibility Meter. Sub-samples were ground using a pestle and mortar, to achieve a homogeneous sample, and sieved at 0.25mm to remove any large clasts prior to analysis. X-ray Fluorescence (XRF) was undertaken on ground samples on a Panalytical Epsilon 3-XL at the School of Geography at the University of Nottingham with resulting spectra analysed to give values for the major oxides and elements MgO, Al_2O_3 , SiO_2 , K_2O , CaO , Ti , Fe_2O_3 and Sr . Particle Size Analysis was undertaken using a Coulter LS 200 Laser Granulometer after samples had been sieved at 1.4mm and disaggregated using a weak sodium hexametaphosphate solution. The GRADISTAT software package (Blott and Pye, 2001) was used to analyse this data.

3. Results

3.1 Sedimentology

The field descriptions of the five off-site sedimentary sections nearest to the site can be found below and their relative locations are shown in Figures 2 and 4. Section 3 was dug into the wadi south of the site and is not described here. In general, there are two major sedimentary units around the site 1) a series of pale coloured fine silts that make up the terrace on which the site sits, and 2) a series of reddish-brown, silts, sands and gravels (with clasts of flint) that are found in the wadi running to the south of the site.

KHIV Section 1 (31 43' 27.1" N; 36 27' 05.4" E)

0 – 21 cm Light Red (10YR 7/6) silty sand with occasional roots. At base (16 – 21cm) large (2-5cm) flint clasts, some of which lie flat on the base of the unit.

21 – 26cm Pink (10YR 10/4) sand. Contains carbonate concretions and has secondary ?salt features suggesting soil formation during period of stasis or drying episode. Not laterally continuous over the site – predates an erosional episode prior to or during deposition of unit above.

26 – 30cm Same as the basal 5cm of top unit showing erosional features, rip up clasts into unit below and erosional surface on the upper contact.

30 – 62cm Light greenish grey (Gley 7/5GY) silty clay with very occasional large (>10cm) flint clasts. OSL sample GL11035 was taken from the top of this unit.

KHIV Section 2 (31 43' 27.8" N; 36 27' 05.6" E)

0 – 30cm Weathering surface and drape

30 – 73cm Pinkish white (10YR 8/2) homogenous silt. OSL sample GL11036 taken from 56cm depth.

KHIV Section 4 (31 43' 22.5" N; 36 27' 13.0" E)

0-12cm Very pale brown (10YR 7/4) silty fine sand containing small (1-3cm) clasts of flint and stone with roots to surface.

12-27cm Brownish yellow (10YR 6/6) fine sand with 2-5cm scale flint clasts. OSL sample GL11037 taken from this unit.

27-82 cm Light yellowish brown (10YR 6/4), but mottled, clayey silt with numerous fine roots and some larger roots.

232 *KHIV Section 5 (31 43'21.6" N; 36 27'21.0" E)*

233

234 0-10cm Weathering surface

235 10-92cm Very pale brown (10YR 8/2) silt with abundant root holes. OSL samples

236 GL11038 and GL11039 taken from 40cm and 80cm respectively.

237 92-112cm Very pale brown (10YR 7/3) sandy silt with occasional small (1-2mm) flint
238 specs.

239 112-137cm Very pale brown (10YR 7/4) silts with common small specs (1-2mm) of flint
240 and charred plant remains or charcoal and occasional large (3-5cm) flints.

241

242 *KHIV Section 6 (31 43' 22.7" N; 36 27' 18.0" E)*

243

244 0-10cm Yellow (10YR 7/6) fine sand containing roots to the surface and plant
245 remains. There is a dry crust on the Wadi surface.

246 10-36cm Light yellowish brown (10YR 6/4) sandy silt containing root holes and organic
247 remains and occasional small (<1cm) flints. OSL sample GL11040 taken from the base of
248 this section.

249 36 – 56cm Very pale brown (10YR 7/3) sand and gravel with large (>10cm) flint clasts.

250 56-111cm Yellowish brown (10YR 5/4) silty clay with occasional large flints.

251

252 *KHIV GeoTrench*

253 *Description based on Locus Summary Sheets*

254

255 0cm Locus 000 Surface

256 0-10cm Locus 001 Disturbed 10cm of slopewash on surface

257 10 – 25cm Locus 002 Red/brown Wadi Silt. OSL sample 11041 taken from this locus.

258 25 – 40cm Locus 003 Brown Clay – possible palaeosol in lake sediments?

259 40 – 43cm Locus 004 Upper White Clay

260 43 – 53cm Locus 005 Brown Clay. OSL sample GL11042 taken from this locus.

261 53 – 55cm Locus 006 Middle White Clay

262 55 – 70cm Locus 007 Brown 'Palaeosol'; looks like level of earliest site occupation given
263 stratigraphy and comparison to sediments in AZ51.

264 70-72cm Locus 008 Lower White Clay looks like the white silt seen towards the base of
265 AZ51 and in Sections 2 and 5

266

267 The sedimentological analyses (Table 1) show distinct differences between the on-site and
268 off-site sediments particularly in terms of magnetic susceptibility and the amounts of calcium

and silica in the sediment. Average matrix grain size (Fig. 5) is slightly coarser on-site, most likely reflecting the anthropogenic origin of some of these sediments.

Given that there is little difference in iron content between the groups of samples it is likely that the substantial differences in magnetic susceptibility values reflect the amount of burning of the sediment onsite (e.g. Morinaga et al., 1999). Analysis of more samples would be required to confirm if the differences between Areas A and B are significant in terms of different intensities of burning throughout these areas, or if certain occupation layers have typically high magnetic susceptibility values. But these preliminary results do suggest this is an area of analysis that may warrant further investigation at the site.

The differences in elemental composition of the sediment, particularly in terms of calcium and silica reflect the visual description of these samples. CaO values from off-site samples are slightly skewed by the carbonate-rich silts of the terrace on which the site sits; samples in sections 1, 2 and 5 have an average of 42.2%. Sieving of these samples for other analyses, and during flotation on-site, reveals that they are rich in ostracod shells and we interpret them as carbonate-enriched wetland sediments, or marl.

3.2 Age-estimates

To provide an absolute chronology for the off-site sediments, where unlike in the site itself charcoal was not preserved, a series of OSL and U-Series age estimates were obtained (Tables 2 and 3). These, along with the radiocarbon chronology of the site, are summarised stratigraphically in Figure 4. Due to low amounts of material suitable for analysis in some OSL samples, some of these age estimates have to be treated with caution or as minimum ages. The lack of material in samples GL-11035, 11039, and 11041 restricted the number of aliquots available for D_e estimation. The latter 2 samples also did not have enough material to allow a dose recovery test, nor did samples 11046 and 11047. Sample GL11043 had significant feldspar contamination, such that this age is a minimum age estimate.

Analytically this leaves seven secure OSL age estimates. In the pale terrace silts we discount samples GL11044 and GL11045 based on stratigraphic reasons. These samples were taken from archaeologically sterile sediments directly below a well-constrained site age (Richter et al., 2013) and therefore cannot be younger than the site. The three other OSL age estimates, GL-11036, 11038 and 11042 give an age of 19 – 23 ka BP for the terrace silts. Although some caution is warranted in the use of this age range as the ‘true’ age of this unit, due to the clearly ‘young’ age estimates of samples GL11044 and GL11045, it is an age

that is supported by 4 of the age estimates from samples with limited datable material, and by the stratigraphic overlap of this unit with the site itself (19,830-18,600 cal. years BP; Richter et al., 2013).

From the second major sedimentary unit (the reddish browns silts, sands and gravels) two OSL age estimates (GL 11037 and 11040) place these deposits in the mid to late Holocene, 5 – 3.5 ka BP. There are no analytically insecure age estimates from this unit, and stratigraphically they sit within the present day wadi, overlying the terrace silts in Section 1.

The individual U-Series ages on the carbonate nodules from the terrace surface gave an average age of $22,740 \pm 920$ years. Individual ages were not corrected for detrital contamination, which $^{230}\text{Th}/^{232}\text{Th}$ ratios show could be important (Table 3; low detrital contamination is usually indicated by high $^{230}\text{Th}/^{232}\text{Th} > 25$; Candy et al. 2005). Although the uncorrected ages of all carbonate concretion subsamples seem highly coherent, an isochron age was calculated to try and take into account this contamination. However, the MSWD (120) and probability of fit (0), given by ISOPLOT as statistical assessment of the fit between the isochron and the original dataset, indicate a probable large degree of scatter around the best-fit isochron such that this age-estimate should be treated with great caution. Failure to obtain a statistically-meaningful isochron can be due to the fact that, although belonging to the same layer, the subsamples did not deposit at exactly the same time or contain different generations of carbonates (Candy et al. 2004); and/or there was more than one source of detrital contaminants. The subsamples also have very similar U-series ratios (Table 3), making it difficult to produce a well-defined isochron (e.g. Dean et al., 2015). This is exemplified by the degree of scatter shown on selected activity ratios (AR) plots (Fig. 6). Although the Rosholt plots, $^{230}\text{Th}/^{232}\text{Th}$ AR versus $^{238}\text{U}/^{232}\text{Th}$ (Rosholt I plot) and $^{234}\text{U}/^{232}\text{Th}$ AR versus $^{238}\text{U}/^{232}\text{Th}$ (Rosholt II plot; Rosholt et al. 1976), emphasise alignment of subsamples, which suggests suitability for the construction of an isochron, the Osmond plots, $^{230}\text{Th}/^{238}\text{U}$ AR versus $^{232}\text{Th}/^{238}\text{U}$ (Osmond I plot) and $^{234}\text{U}/^{238}\text{U}$ AR versus $^{232}\text{Th}/^{238}\text{U}$ (Osmond II plot; Osmond et al. 1970) highlight the clustering of subsamples due to chemical similarities which render them inappropriate for statistically meaningful isochron calculations.

The age of ca. 22,500-23,500 years given by both the individual dates and the isochron date should therefore be considered as a maximum age of the sample (since it cannot be properly corrected for initial detritalTh). This fits with the other chronological and stratigraphic controls on the site, as we presume these nodules formed during, or after the silts in which they sit (e.g. Rowe & Maher 2000) i.e. after c. 19 ka BP.

4. Discussion

The detailed mapping and absolute dating of the sediments surrounding KHIV allows us to reconstruct the environmental changes at the site for various time windows over the last 23,000 years.

The spatial extent, duration and type of water body that deposited the pale terrace clays and silts at KHIV are difficult to establish. The present day extent of the marl terrace is clear from satellite imagery (Fig. 3) but as Garrard et al. (1985) noted there is no clear natural barrier to form a lake in this point on the wadi, and there are no shorelines evident against the limestone bedrock on the northern edge of the wadi. A more recent drainage pattern is now superimposed on the wadi, cutting the marl terrace between the site and section 5, and the main drainage channel of the WadiKharaneh to the north of the site may have eroded any remaining shoreline evidence.

The lack of distinct shoreline and other morphological features of the marl, such as the apparent parallel nature of the sedimentary units to the wadi floor, are similar to those defining ground-water discharge (GWD) deposits (Pigati et al., 2014). The sediments, especially those described in sections 2 and 5 often resemble those described as 'Wetland Marl' by Pigati et al. (2014) i.e. massive to blocky, which they interpret as forming in shallow wetlands, or in marshy areas. Of note at Kharaneh though is the massive nature of some of the marl, particularly in Section 2, suggesting there was little vegetation growing at the site of deposition. This suggests that the Kharanehwetland, at least at times, held substantial amounts of water and may have had open water areas.

Interpretation of this water body as being a GWD deposit is hard to envisage given the main Azraqaquifers today are at least 100m below Kharaneh (e.g. Al-Kharabsheh, 2000). However, in times of more effective precipitation (see further discussion below) it is possible there was a localised, shallow, groundwater source at this location. Surface water recharge of this wetland may also have been possible. The marl terrace and the site sit in a particularly wide section of the main wadi channel, constrained by the limestone bedrock wadi edge to the north and the flint pavement (D in Figure 3) to the south. The full depth of the 'basin' in which the Pleistocene sediments of Kharaneh sit is unknown, but based on current topography this is a section of the wadi where flowing surface water could have slowed down and pooled, particularly in an area already rich in wetland vegetation. The recent digging of a dam near KHIV (clearly visible in Fig. 3) has shown that winter rains draining through WadiKharaneh today can last well into the summer months given sufficient

storage capacity. Given the spatial extent of the marl terrace it is likely that the Late Pleistocene wetland that produced these sediments would have been in the order of 50 times larger than this dam, at least at its maximum extent.

The marl terrace, and therefore the wetland which deposited it, dates to between 23 and 19 kaBP, based on the chronological discussion above. Today, there is a slight stratigraphic overlap between the top of sections 2 and 5, and the occupation levels of AZ51 on-site, given the regional topographic gradient (Figure 4). However, it is likely that this terrace was higher in the past. Besancon et al. (1989) describe carbonate concretions at the base of a 30cm silt layer, we observe these nodules (from which the U-Series age estimates were produced) at or near the surface today, suggesting some substantial deflation in the ~30 years between our surveys.

Under the western and northern areas of the site itself (Area A and AZ51), the wetland deposits are ostracod-rich, carbonate-concreted greenish marls, similar to those seen in sections 2 and 5, and are interstratified with the earliest Early EP occupations. Under the eastern portion of the site (Area B), archaeologically sterile, tan-coloured clays with little visible carbonate form an abrupt boundary (with no visible mixing) with the overlying occupational deposits (Maher et al., this volume). Given the subtle differences in the wetland facies observed on and around the site and their stratigraphic overlap with the site itself, and the microfossils observed during our initial analyses presented here (i.e. ostracods; diatoms are also preserved, K. Mills pers. com.) more detailed analysis of these wetland sediments are planned to tease out the detail of environmental change recorded here through the late Pleistocene.

Following the marl deposition Besancon et al. (1989) and Garrard et al. (1985) describe a silty loam (with carbonate concretions at the base) which today appears to have largely been deflated. Garrard et al. (1985) suggest these were loess deposits that, given our chronological data, were deposited at some point post-19 ka BP and would suggest substantial drying of the local environment. It's possible that these loess deposits are the same as those found in Locus 2 of the Geotrench (with a cautious age estimate of 15 ± 1 ka BP; GL11041) but we are not able to link them together directly. The next depositional event related to the site is the Holocene fill identified in the minor wadis that make up the present day drainage pattern, dating to around 4 ka BP. This points to a substantial erosional phase of the marl terrace at some time between 19ka and 4ka BP.

4.1 A suitable site for occupation?

Given the location of the site, above much of the marl deposition, and also within the southern limits of the proposed maximum extent of the wetland(Fig. 3), it is unlikely that the most extensive Pleistocene water body still existed at Kharaneh at the time of the first site occupation around 20,000 years ago. However, as reported above, it seems likely that water did still exist at the site to some degree when it was first occupied, at least on a seasonal basis. Given the site's environmental history prior to occupation as documented here,Kharaneh IV would likely have appeared anoptimal location within a resource-rich environment in which to set up camp. The sustained occupation of the site suggests, despite limited sedimentary evidence post 19 ka BP, these resources were available for some time, at least 1200 years.

Jones and Richter (2011) show that the centralAzraq oasis was also a well-watered locale at this time and yet there is no large aggregation site apparent there. Archaeological evidence suggests that groups using different sets of lithic technology and with ties to either the west or the southern and northern Levant occupied the Azraq Basin during the Early Epipalaeolithic (Richter et al., 2011; Maher et al., this volume). It is possible that social barriers prevented the establishment of a large basecamp-style aggregation site in the oasis itself at that time. The AzraqOasis may have fallen in between territories of different social groups of hunter-gatherers making the establishment of a large site here socially unacceptable. This idea is supported by the fact that the only other large aggregation site in the Azraq Basin, Jilat 6, is characterised by a very different set of lithic industries compared to KHIV, whereas the lithic assemblages recovered from Ayn Qasiyya, a smaller site,have parallels with both the KHIV and Jilat 6 lithic assemblages. At the same time, it is also possible that the oasis may not have been suitable for long-term aggregated settlement due to other factors, such asthe presence of large predators. Given the long history of archaeological survey in and around the oasis it is unlikely that a site of the magnitude of Jilat 6 or KHIV has been missed.

Unfortunately there is no local sedimentary evidence from which to reconstruct the environment through most of the occupation of KHIV, or to point to reasons for eventual site abandonment. Such environmental information must come from ongoing work from the site itself. The now largely deflated loess deposits described by Garrard et al. (1985) and Besancon et al. (1989) does suggest a drier period following the wetlanddeposits that overlap with the site but there are nostratigraphically secure absolute dates to confirm if these were deposited during the site occupation, or following abandonment.

4.2 Comparison to regional palaeoenvironmental records

High lake levels during the Late Pleistocene are reported from across the wider eastern Mediterranean region, with water bodies substantially larger than those found today, such as Lake Lisan (e.g. Torfstein et al., 2013), Lake Van (Çağatay et al., 2014) and in the Konya plain (e.g. Roberts, 1983). A combination of increased precipitation and/or reduced evaporation is likely to have increased the potential (compared to present day conditions) for standing water to remain, where geomorphological conditions allowed. Both Lake Lisan and Konya had significant falls in lake levels ~ 21 ka BP and the deposits at Kharaneh IV would fit this pattern with the maximum extent of water at the site occurring before site occupation around 20 ka BP, and subsequent drying afterwards.

Evidence from other sites in the Azraq Basin would also suggest that the period of most positive water balance in the basin occurred shortly prior to 20 ka BP. Garrard et al. (1988, 1994) and Garrard and Byrd (2013) interpret the sediments of Uweynid 14 (23.4 – 21.4 kcal BP; Richter et al., 2013) as being deposited during a period of relatively high water table and identify a 'humid' phase in the Wadi Jilat around 23 kcal BP (19,000 uncalibrated radiocarbon years BP). The timing of both these events would fit with the absolute dating of the Kharaneh marls. Organic marsh deposits are well established in the central oasis at Ayn Qasiyya by 24 ka BP as water levels fell from a more extensive open water body, although locally open water conditions there continued until 16 ka BP (Jones and Richter, 2011).

There is a lack of continuous post-Last Glacial Maximum sediments in the wider basin that make reconstructing environmental changes through the last glacial-interglacial transition and the early Holocene here difficult. For example in the central oasis there is a sedimentary hiatus at Ayn Qasiyya between 16 and 10.5 ka BP (Jones and Richter, 2011); we cannot therefore place events such as the net erosive period at KHIV between 19 and 4 ka BP with any better resolution. Identifying how environments in the Azraq Basin changed through this important transition remains a particular challenge of work in the region.

5. Conclusions

The Kharaneh wetland was likely a well-known landscape feature for Early Epipalaeolithic occupants of the Azraq Basin. As elsewhere in the region, a relatively positive hydroclimatic balance existed c. 23 ka BP. Water balance has not been as positive in the region since, having already begun to decline by the time of occupation at KHIV. With the central oasis providing persistent water and associated floral and faunal resources throughout this time period, KHIV and Jilat 6 additionally suggest the end of the Pleistocene was a prime time for people to thrive in the Azraq Basin, with a c. 1000 year window of rich

environmental resources that were substantially exploited by Early and Middle EP communities.

Acknowledgements

We thank the Department of Antiquities of Jordan and its Director General, our DoA reps for the field seasons at Kharanehand our DoA collaborator Ahmad Lash. We thank all the field crews of 2007-2015, in particular Samantha Allcock, and the community of Azraq for their continued support. This work was funded by grants from the Arts and Humanities Research Council of Britain, Council for British Research in the Levant, American Centre for Oriental Research, University of California, Berkeley, the Wenner-Gren Foundation, the University of Nottingham, and the Fragmented Heritage Project at the University of Bradford. Phil Toms is thanked for his advice on the interpretation of the OSL age estimates. This manuscript was written while MJ was a visiting Fellow in the School of Geography, Planning and Environmental Management at the University of Queensland. We thank Jason Rech and an anonymous reviewer for comments which improved this manuscript.

References

- Al-Juaidi, A. E., Kaluarachchi, J. J., & Mousa, A. I. (2014). Hydrologic-Economic Model for Sustainable Water Resources Management in a Coastal Aquifer. *Journal of Hydrologic Engineering*, 19(11).
- Al-Kharabsheh, A. (2000). Ground-water modelling and long-term management of the Azraq basin as an example of arid area conditions (Jordan). *Journal of arid environments*, 44(2), 143-153.
- Ames, Chris, April Nowell, Carlos Cordova, James Pokines and Michael Bisson. 2014. The Druze Marsh Paleolandscape: A Geoarchaeological Approach to Open-Air Paleolithic Sites. *Quaternary International* 331, 60-73.
- Berger, G.W., Mulhern, P.J. and Huntley, D.J. (1980). Isolation of silt-sized quartz from sediments. *Ancient TL*, 11, 147-152.
- Berndtsson, R., Jebbari, S., Hashemi, H., & Wessels, J. (2014). Traditional water management techniques—do they have a role in post Arab Spring Middle East? *Hydrological Sciences Journal*.
- Besançon, J., Geyer, B., Sanlaville, P., 1989. Contribution to the study of the geomorphology of the Azraq Basin, Jordan. In: Copeland, L., Hours, F. (Eds.), *The Hammer on the Rock: Studies in the Early Palaeolithic of Azraq*. British Archaeological Reports International Series 540, Oxford, pp. 7-63.
- Betts, A.V.G., 1998. *The Harra and the Hamad. Excavations and Surveys in Eastern Jordan*, Volume 1. Sheffield Academic Press, Sheffield.
- Black S., Robinson S., Fitton R., Goodship R., and Rambeau C.M.C. 2011. Palaeoenvironmental and limnological reconstruction of Lake Lisan and the Dead Sea. In *Water Life and Civilisation: Climate, Environment and Society in the Jordan Valley*. International Hydrology Series. S. Mithen and E. Black (eds), Cambridge: Cambridge University Press
- Blockley, S. P. E., & Pinhasi, R. (2011). A revised chronology for the adoption of agriculture in the Southern Levant and the role of Lateglacial climatic change. *Quaternary Science Reviews*, 30(1), 98-108.
- Blott, S.J. and Pye, K. (2001) GRADISTAT: a grain size distribution and statistics package for the analysis of unconsolidated sediments. *Earth Surface Processes and Landforms* 26, 1237-1248.
- Çağatay, M. N., Öğretmen, N., Damcı, E., Stockhecke, M., Sancar, Ü., Eriş, K. K., & Özeren, S. (2014). Lake level and climate records of the last 90ka from the Northern Basin of Lake Van, eastern Turkey. *Quaternary Science Reviews*, 104, 97-116.
- Candy, I., Black, S. and Sellwood, B.W. 2005. U-series Isochron Dating of Immature and Mature Calcretes as a Basis for Constructing Quaternary Landform Chronologies for the

542 Sorbas Basin, Southeast Spain. *Quaternary Research*, 64 pp. 100-111.

543 Candy, I., Black, S., Sellwood, B.W., 2004a. Quantifying timescales of pedogenic calcrete
544 formation using U-series disequilibria. *Sedimentary Geology* 170, 177–187.

545 Copeland, L., and F. Hours. Editor. 1989. *The Hammer on the Rock: Studies in the Early*
546 *Palaeolithic of Azraq, Jordan*. Oxford: B.A.R. International Series 540.

547 Cordova, Carlos, April Nowell, Michael Bisson, Christopher Ames, James Pokines, Melanie
548 Chang and Maysoon al-Nahar. 2013. Glacial and Interglacial Desert Refugia and the Middle
549 Paleolithic of the Azraq Basin, Jordan. *Quaternary International* 300, 194-210

550 Dean, J.R., Jones M.D., Leng, M.J., Noble, S.R., Metcalfe, S.E., Sloane H.J., Sahy, D.,
551 Eastwood, W.J. and Roberts, C.N. 2015. Eastern Mediterranean hydroclimate over the late
552 glacial and Holocene, reconstructed from the sediments of Nar Gölü, central Turkey, using
553 stable isotopes and carbonate mineralogy. *Quaternary Science Reviews* 124, 290-295.

554 Enzel, Y., Amit, R., Dayan, U., Crouvi, O., Kahana, R., Ziv, B., & Sharon, D. (2008). The
555 climatic and physiographic controls of the eastern Mediterranean over the late Pleistocene
556 climates in the southern Levant and its neighboring deserts. *Global and Planetary*
557 *Change*, 60(3), 165-192.

558 Field, H. 1960. *North Arabian Desert Archaeological Survey, 1925-1950. Papers of the*
559 *Peabody Museum*. Cambridge: Peabody Museum, Harvard University.

560 Galbraith, R. F., Roberts, R. G., Laslett, G. M., Yoshida, H. and Olley, J. M. (1999) Optical
561 dating of single and multiple grains of quartz from Jinmium rock shelter (northern Australia):
562 Part I, Experimental design and statistical models. *Archaeometry*, 41, 339-364.

563 Garrard, A. N. and Byrd, B. 2013. *Beyond the Fertile Crescent: Late Palaeolithic and*
564 *Neolithic communities of the Jordanian steppe. The Azraq Basin Project Volume 1: Project*
565 *background and the Late Palaeolithic (geological context and technology)*. Oxford: Oxbow.

566 Garrard, A. N., 1998. Environment and cultural adaptations in the Azraq Basin: 24,000 -
567 7,000 B.P. In: Henry, D.O., (Ed.) *The Prehistory of Jordan*. British Archaeological Reports
568 International Series 705, Oxford, pp. 139-48

569 Garrard, A. N., Baird, D., Colledge, S., Martin, L., Wright, K., 1994. Prehistoric Environment
570 and Settlement in the Azraq Basin: an interim report on the 1987 and 1988 Excavation
571 Season. *Levant* 26, 73-109.

572 Garrard, A., Byrd, B., Harvey, P., & Hivernel, F. (1985). Prehistoric environment and
573 settlement in the Azraq Basin. A report on the 1982 survey season. *Levant*, 17(1), 1-28.

574 Heiri, O., Lotter, A. F., & Lemcke, G. (2001). Loss on ignition as a method for estimating
575 organic and carbonate content in sediments: reproducibility and comparability of
576 results. *Journal of paleolimnology*, 25(1), 101-110.

577 IPCC, 2014: Summary for policymakers. In: *Climate Change 2014: Impacts, Adaptation, and*
578 *Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the*

579 Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Field, C.B.,
580 V.R. Barros, D.J. Dokken, K.J. Mach, M.D. Mastrandrea, T.E. Bilir, M. Chatterjee, K.L. Ebi,
581 Y.O. Estrada, R.C. Genova, B. Girma, E.S. Kissel, A.N. Levy, S. MacCracken, P.R.
582 Mastrandrea, and L.L. White (eds.)]. Cambridge University Press, Cambridge, United
583 Kingdom and New York, NY, USA, pp. 1-32

584 Jackson, M.L., Sayin, M. and Clayton, R.N. (1976). Hexafluorosilicic acid regent modification
585 for quartz isolation. *Soil Science Society of America Journal*, 40, 958-960.

586 Jones, J. R. (2012). Using gazelle dental cementum studies to explore seasonality and
587 mobility patterns of the Early-Middle Epipalaeolithic Azraq Basin, Jordan. *Quaternary*
588 *International*, 252, 195-201.

589 Jones, M. and T. Richter. 2011. Palaeoclimatic and archaeological implications of
590 Pleistocene and Holocene environments in Azraq, Jordan. *Quaternary Research* Vol. 76 (3),
591 363-372

592 Ludwig, K.R. 2001. ISOPLOT/Ex rev. 2.49, United States Geological Survey.

593 Maher L.A. Macdonald, D.A., Allentuck, A., Martin, L., Spyrou, A and Jones, M.D. *this*
594 *volume* Occupying Wide Open Spaces? Late Pleistocene Hunter-Gatherer Activities in the
595 Eastern Levant

596 Maher, L. (In review). Late Quaternary Refugia, Aggregations and Palaeoenvironments in
597 the Azraq Basin. In *Quaternary Environments, Climate Change and Humans in the*
598 *Levant*. Cambridge University Press: Cambridge.

599 Maher, L. A., Banning, E. B., & Chazan, M. (2011). Oasis or mirage? Assessing the role of
600 abrupt climate change in the prehistory of the southern Levant. *Cambridge Archaeological*
601 *Journal*, 21(1), 1-29.

602 Maher, L., Richter, T., Jones, M., Stock, J.T., 2011b. The Epipalaeolithic Foragers in Azraq
603 Project: Prehistoric Landscape Change in the Azraq Basin, Eastern Jordan. CBRL Bulletin 6,
604 21-27.

605 Maher, L.A., Richter, T., Macdonald, D., Jones, M.D., Martin, L., Stock, J.T., 2012. Twenty
606 thousand-year-old huts at a hunter-gatherer settlement in eastern Jordan. PloS one 7,
607 e31447.

608 Martin, L., Edwards, Y., & Garrard, A. (2010). Hunting practices at an eastern Jordanian
609 Epipalaeolithic aggregation site: the case of Kharaneh IV. *Levant*, 42(2), 107-135.

610 Morinaga, H., Inokuchi, H., Yamashita, H., Ono, A., & Inada, T. (1999). Magnetic detection of
611 heated soils at Paleolithic sites in Japan. *Geoarchaeology*, 14(5), 377-399.

612 Muheisen M (1988) The Epipalaeolithic phases of Kharaneh IV. In: Garrard A, Gebel H,
613 editors. The Prehistory of Jordan The State of Research in 1986. Oxford: British
614 Archaeological Reports 396. pp. 353-367.

615 Muheisen M (1988) The Epipalaeolithic phases of Kharaneh IV. In: Garrard A, Gebel H,
616 editors. The Prehistory of Jordan The State of Research in 1986. Oxford: British
617 Archaeological Reports 396. pp. 353-367.

618 Murray, A.S. and Wintle, A.G. (2000) Luminescence dating of quartz using an improved
619 single-aliquot regenerative-dose protocol. *Radiation Measurements*, 32, 57-73.

620 Murray, A.S. and Wintle, A.G. (2003) The single aliquot regenerative dose protocol: potential
621 for improvements in reliability. *Radiation Measurements*, 37, 377-381.

622 Osmond, J. K., J. P. May, and W. F. Tanner (1970), Age of the Cape Kennedy Barrier-and-
623 Lagoon Complex, Journal of Geophysical Research, 75, 469-479.

624 Pigati, J.S., Rech, J.A., Quade, J. and Bright, J. (2014). Desert wetlands in the geologic
625 record. *Earth-Science Reviews*, 132, 67-81.

626 Prescott, J.R. and Hutton, J.T. (1994) Cosmic ray contributions to dose rates for
627 luminescence and ESR dating: large depths and long-term time variations. *Radiation*
628 *Measurements*, 23, 497-500.

629 Rambeau C.M.C., Finlayson B., Smith S., Black S., Inglis R. and Robinson S. 2011.
630 Palaeoenvironmental reconstruction at Beidha, southern Jordan (c. 18,000-8,500 BP):
631 implications for human occupation during the Natufian and Pre-Pottery Neolithic. In *Water*
632 *Life and Civilisation: Climate, Environment and Society in the Jordan Valley*. International
633 Hydrology Series. S. Mithen and E. Black (eds), Cambridge: Cambridge University Press

634 Ramsey, M.N., Jones, M.D., Richter, T. and Rosen, A. *in press* Modifying the marsh:
635 Evaluating Early Epipaleolithic hunter-gatherer impacts in the Azraq wetland, Jordan *The*
636 *Holocene*

637 Richter, T, Garrard, A, Allcock, S & Maher, L 2011, 'Interaction Before Agriculture:
638 Exchanging Material and Sharing Knowledge in the Final Pleistocene Levant' *Cambridge*
639 *Archaeological Journal*, vol 21, no. 1, pp. 95-114.

640 Richter, T., Maher, L.A., Garrard, A.N., Edinborough, K., Jones, M.D., Stock, J.T., 2013.
641 Epipalaeolithic settlement dynamics in southwest Asia: new radiocarbon evidence from the
642 Azraq Basin. *Journal of Quaternary Science* 28, 467-479.

643 Roberts, N. (1983). Age, palaeoenvironments, and climatic significance of late Pleistocene
644 Konya Lake, Turkey. *Quaternary Research*, 19(2), 154-171.

645 Robinson, S. A., Black, S., Sellwood, B. W., & Valdes, P. J. (2006). A review of
646 palaeoclimates and palaeoenvironments in the Levant and Eastern Mediterranean from
647 25,000 to 5000 years BP: setting the environmental background for the evolution of human
648 civilisation. *Quaternary Science Reviews*, 25(13), 1517-1541.

649 Rollefson, G., Schnurrenberger, D., Quintero, L., Watson, R.P., Low, R., 1997. Ain Soda and
650 'Ayn Qasiya: New late pleistocene and early Holocene sites in the Azraq Shishan area,

651 eastern Jordan. In: Gebel, H.G.K., Kafafi, Z., Rollefson, G.O. (Eds.) The prehistory of Jordan
652 II. Perspectives from 1997. ex oriente, Berlin, 45-58.

653 Rosen, A. M. (2007). *Civilizing climate: social responses to climate change in the ancient*
654 *Near East*. Rowman Altamira.

655 Rosen, Arlene M., and Isabel Rivera-Collazo 2012 Climate Change, Adaptive Cycles, and
656 the Persistence of Foraging Economies during the Late Pleistocene/Holocene Transition in
657 the Levant. *Proceedings of the National Academy of Sciences* 109(10): 3640–3645.

658 Rosholt, J.N. (1976) ²³⁰Th/U dating of travertine and caliche rinds. The Geological Society of
659 America, Abstracts and Program, 8: 1076 p.

660 Rowe P.J., Maher, B.A. (2000). 'Cold' stage formation of calcrete nodules in the Chinese
661 Loess Plateau: evidence from U-series dating and stable isotope analysis.
662 *Palaeogeography, Palaeoclimatology, Palaeoecology* 157, 109–125

663 Ruddiman, W. F., Ellis, E. C., Kaplan, J. O., & Fuller, D. Q. (2015). Defining the epoch we
664 live in. *Science*, 348(6230), 38-39.

665 Torfstein, A., Goldstein, S. L., Stein, M., & Enzel, Y.
666 (2013). Impacts of abrupt climate changes in the Levant from Last Glacial Dead Sea
levels. *Quaternary Science Reviews*, 69, 1-7.

Figure and Table Captions

Figure 1 Extent of the Azraq basin (dashed line) showing major wadis (solid lines) and playa (hashed areas). The major archaeological sites discussed in the text are shown. Shading depicts 250m contour intervals (masl). The sites of Kharaneh IV and Ayn Qassiyra are also shown on the regional map for context.

Figure 2 Detailed topography of the WadiKharaneh around the site of Kharaneh IV (here marked by the thick black line). The locations of the off-site sedimentary sections as described in the text are shown. The dotted line marks the transect described in Figure 4.

Figure 3 Annotated satellite image of the site and surrounding area from Google Earth. The dotted line represents the maximum extent of the Kharaneh wetland, as defined by the bedrock topography and distribution of marl sediments.

Figure 4Relative distribution of off-site stratigraphy and occupational horizons of Kharaneh IV. All age estimates associated withthese sections are shown (bar one 'rejected' OSL date for the Geo Trench; GL11041); age estimates in bold are used in the final interpretations. The location of OSL samples are marked by circles (see Table 1 for details), the date of the present-day terrace surface was obtained using U-Series techniques (Table 2) and the date for the occupation horizons comes from Bayesian modelling of 13 radiocarbon age estimates (Richter et al., 2013). Note; for clarity the location of Section 4 has been moved, its actual location marked by the dotted outline. A full description of the sub-units of each section can be found in the main text.

Figure 5Particle size summaries for analysis of sediments taken from on- and off-site sections at KIV.

Figure 6Activity Ratio bi-plots from the U-Series analysis of carbonate nodules from near Kharaneh IV following Rosholt et al. (1976) and Osmond et al. (1970)

Table 1Data summary from sediment samples from Kharaneh IV and surrounding sediments. Mean values \pm 1 standard deviation are shown.

Table 2Dose rate (D_r), Equivalent Dose (D_e) and Age data from Kharaneh IV OSL samples. Further discussion of the samples listed as having limited datable material or significant feldspar contamination can be found in the main text.

704

705 **Table 3** Uranium/Thorium age for sample KAL-IV. The isochron age is calculated using a
706 series of subsamples (1-5). Uncorrected U/Th ages for each subsample are given in italics.
707 Average uncertainties (SDs) on U and Th concentrations are calculated from all data
708 measured during the same batch and are 0.45% and 0.67% respectively.

709

710

Figure 1




Figure 2





Figure 3




Figure 4




Figure 5




Figure 6




Table 1

Table 1 Data summary from sediment samples from Kharaneh IV and surrounding sediments. Mean values \pm 1 standard deviation are shown.

Samples locations	Carbon content % weight loss at 550°C	Carbonate content % weight loss at 925°C	Magnetic Susceptibility	MgO %	Al ₂ O ₃ %	SiO ₂ %	K ₂ O %	CaO %	Ti %	Fe ₂ O ₃ %	Sr %
Area A (n=11)	9.0 \pm 3.5	10.7 \pm 2.6	440.7 \pm 100.8	6.2 \pm 1.4	9.6 \pm 1.0	46.8 \pm 3.4	1.2 \pm 0.1	26.0 \pm 3.7	0.7 \pm 0.1	7.3 \pm 1.0	0.3 \pm 0.1
Area B (n=24)	8.5 \pm 2.3	11.3 \pm 2.8	224.4 \pm 157.6	5.1 \pm 0.7	10.7 \pm 1.2	46.0 \pm 6.0	1.4 \pm 0.3	26.2 \pm 7.4	0.8 \pm 0.1	9.2 \pm 1.3	0.2 \pm 0.1
Off-site (n=17)	7.8 \pm 2.3	13.7 \pm 5.3	74.5 \pm 15.9	3.8 \pm 0.4	9.8 \pm 1.5	39.9 \pm 5.5	1.0 \pm 0.4	35.8 \pm 8.3	0.7 \pm 0.1	8.2 \pm 1.1	0.3 \pm 0.2

Table 2

Table 2 Dose rate (D_r), Equivalent Dose (D_e) and Age data from Kharaneh IV OSL samples. Further discussion of the samples listed as having limited datable material or significant feldspar contamination can be found in the main text.

Field code	Lab code	K %	Th %	U %	αD_r Gy.k ^a ⁻¹	βD_r Gy.k ^a ⁻¹	γD_r Gy.k ^a ⁻¹	Cosmic D_r Gy.k ^a ⁻¹	Total D_r Gy.k ^a ⁻¹	D_e Gy	Age (ka)	Comment
Section 1	GL11035	0.83 ± 0.04	5.42 ± 0.42	1.85 ± 0.1	0.35 ± 0.02	1.02 ± 0.05	0.64 ± 0.03	0.21 ± 0.02	2.23 ± 0.07	39.4 ± 7.2	18 ± 3	Limited datable material
Section 2	GL11036	1.07 ± 0.05	6.22 ± 0.44	2.02 ± 0.11	0.39 ± 0.03	1.25 ± 0.06	0.76 ± 0.04	0.19 ± 0.02	2.59 ± 0.08	51.4 ± 2.7	20 ± 1	
Section 4	GL11037	0.71 ± 0.04	4.44 ± 0.40	2.67 ± 0.13	0.41 ± 0.03	1.03 ± 0.05	0.67 ± 0.03	0.22 ± 0.03	2.33 ± 0.07	10.5 ± 0.8	4.5 ± 0.4	
Section 5 40cm	GL11038	0.84 ± 0.04	5.25 ± 0.38	1.75 ± 0.10	0.33 ± 0.02	1.00 ± 0.05	0.62 ± 0.03	0.16 ± 0.02	2.13 ± 0.07	44.8 ± 2.7	21 ± 1	
Section 5 80cm	GL11039	0.69 ± 0.04	3.83 ± 0.36	1.49 ± 0.09	0.27 ± 0.02	0.84 ± 0.04	0.51 ± 0.03	0.17 ± 0.02	1.78 ± 0.06	34.6 ± 2.3	19 ± 1	Limited datable material
Section 6	GL11040	0.97 ± 0.05	6.76 ± 0.52	2.23 ± 0.11	0.42 ± 0.03	1.21 ± 0.06	0.77 ± 0.04	0.22 ± 0.02	2.62 ± 0.08	10.0 ± 0.7	3.8 ± 0.3	
Geo B OSL 1	GL11041	0.96 ± 0.05	6.06 ± 0.44	2.10 ± 0.11	0.38 ± 0.03	1.15 ± 0.06	0.72 ± 0.04	0.22 ± 0.02	2.47 ± 0.08	37.5 ± 1.5	15 ± 1	Limited datable material
GEO B OSL 2	GL11042	1.14 ± 0.06	6.81 ± 0.46	2.19 ± 0.11	0.42 ± 0.03	1.33 ± 0.07	0.81 ± 0.04	0.21 ± 0.02	2.77 ± 0.09	61.1 ± 3.4	22 ± 1	
AZ51 OSL	GL11043	0.88 ± 0.05	5.79 ± 0.43	1.97 ± 0.10	0.37 ± 0.02	1.08 ± 0.06	0.68 ± 0.03	0.18 ± 0.02	2.32 ± 0.07	30.6 ± 1.5	13 ± 1	Significant feldspar contamination
R/S2/60	GL11044	1.12 ± 0.05	6.46 ± 0.45	1.91 ± 0.10	0.38 ± 0.03	1.27 ± 0.07	0.76 ± 0.04	0.18 ± 0.02	2.59 ± 0.08	36.6 ± 2.0	14 ± 1	
R/S2/60	GL11045	1.02 ± 0.05	5.49 ± 0.42	1.92 ± 0.10	0.36 ± 0.02	1.19 ± 0.06	0.71 ± 0.03	0.19 ± 0.02	2.45 ± 0.08	41.5 ± 2.2	17 ± 1	
BS58 OSL 2	GL11046	0.84 ± 0.05	5.04 ± 0.39	1.83 ± 0.10	0.32 ± 0.02	0.99 ± 0.06	0.61 ± 0.03	0.18 ± 0.02	2.11 ± 0.07	44.6 ± 2.1	21 ± 1	Limited datable material
As42 OSL 4	GL11047	0.67 ± 0.04	4.24 ± 0.4	2.07 ± 0.11	0.34 ± 0.02	0.90 ± 0.05	0.57 ± 0.03	0.17 ± 0.02	1.99 ± 0.06	38.4 ± 1.5	19 ± 1	Limited datable material

Table 3

Table 3 Uranium/Thorium age for sample KAL-IV. The isochron age is calculated using a series of subsamples (1-5). Uncorrected U/Th ages for each subsample are given in italics. Average uncertainties (SDs) on U and Th concentrations are calculated from all data measured during the same batch and are 0.45% and 0.67% respectively.

Sub-sample	U ($\mu\text{g/kg}$)	Th ($\mu\text{g/kg}$)	$^{234}\text{U}/^{238}\text{U}$	$^{230}\text{Th}/^{238}\text{U}$	$^{230}\text{Th}/^{232}\text{Th}$	Age (y.BP)
KAL-IV-1	8820	2412	1.150 ± 0.011	0.187 ± 0.001	2.100 ± 0.015	<i>22509</i> <i>± 910</i>
KAL-IV-2	9832	2848	1.153 ± 0.006	0.190 ± 0.000	2.008 ± 0.007	<i>22827</i> <i>± 922</i>
KAL-IV-3	12102	2256	1.185 ± 0.008	0.191 ± 0.001	3.144 ± 0.015	<i>23016</i> <i>± 930</i>
KAL-IV-4	10018	2955	1.099 \pm 0.008	0.188 ± 0.001	1.956 ± 0.030	<i>22628</i> <i>± 914</i>
KAL-IV-5	7839	1768	1.021 \pm 0.043	0.189 ± 0.004	2.567 ± 0.059	<i>22705</i> <i>± 918</i>
Isochron Age (yrs B.P.)						23560 \pm 1247